



May 21st, 5:00 PM - 5:20 PM

Modelling the polarization, migration and neuromast deposition in the zebrafish posterior lateral line system

Hildur Knutsdottir

University of British Columbia, hildur@math.ubc.ca

Follow this and additional works at: <http://scholarscompass.vcu.edu/bamm>

 Part of the [Developmental Biology Commons](#), [Ordinary Differential Equations and Applied Dynamics Commons](#), and the [Partial Differential Equations Commons](#)

<http://scholarscompass.vcu.edu/bamm/2016/May21/54>

This Event is brought to you for free and open access by the Dept. of Mathematics and Applied Mathematics at VCU Scholars Compass. It has been accepted for inclusion in Biology and Medicine Through Mathematics Conference by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

MODELLING THE POLARIZATION, MIGRATION AND NEUROMAST DEPOSITION IN THE ZEBRAFISH POSTERIOR LATERAL LINE SYSTEM

Hildur Knútsdóttir¹, Cole Zmurchok¹, Dhananjay Bhaskar¹, Eiríkur Pálsson², Leah Edelstein-Keshet¹

¹ Mathematical Biology, University of British Columbia, Vancouver, BC, Canada,

² Department of Biology, Simon Fraser University, Burnaby, British Columbia, Canada.

hildur@math.ubc.ca

Abstract

Collective cell migration plays an important role in a variety of biological systems, such as embryonic development, wound healing and cancer invasion. Here we study a specific example in the zebrafish embryo, where a group of about 100 cells called the posterior lateral line primordium (or PLLP), destined to form sensory structures, migrates from head to tail. Using a 3D deformable cell-based simulation, coupled to reaction-diffusion PDEs for chemical concentrations, we model and simulate the polarization, migration, and neuromast deposition in the PLLP. The model combines realistic cell-cell adhesion and interaction forces, chemotaxis up a gradient of the chemokine SDF-1, and up an FGF ligand gradient to describe the emergence of tissue polarity, as well as cell-cluster shape and speed of collective cell migration. We base the chemical signalling model on receptor-ligand kinetics and mutually inhibitory FGF-Wnt signalling. Our signalling submodel allows predictions to be made for the position of the front-rear boundary in the PLLP, as a function of aggregate parameters such as steady-state receptor expression levels and IC_{50} parameters for the mutually-inhibitory effect. Our 3D model allows an investigation of the role of cell division, chemotaxis, adhesion and other parameters on the shape of the PLLP and demonstrates reasonable behaviour of control as well as mutant phenotypes.